

---

# Clinical Guidance

---

## Paediatric Critical Care: Drowning

### Summary

This guideline is for staff to use when caring for a child following a drowning. It looks at resuscitation, management, investigations and prognostics.

Document Detail	
Document type	Clinical Guideline
Document name	Paediatric Critical Care: Drowning
Document location	Evelina London website
Version	2
Effective from	<i>March 2018</i>
Review date	<i>March 2021</i>
Owner	Jon Lillie, PICU consultant
Author(s)	Jo Perkins, PICU consultant
Approved by, date	Evelina CGG, March 2018
Superseded documents	
Related documents	
Keywords	Evelina, child, Paediatric, intensive care, STRS, PICU Retrieval, Paediatric critical care, Drowning, cardiac arrest, warming
Relevant external law, regulation, standards	
<p>This clinical guideline has been produced by the South Thames Retrieval Service (STRS) at Evelina London for nurses, doctors and ambulance staff to refer to in the emergency care of critically ill children.</p> <p>This guideline represents the views of STRS and was produced after careful consideration of available evidence in conjunction with clinical expertise and experience. The guidance does not override the individual responsibility of healthcare professionals to make decisions appropriate to the circumstances of the individual patient.</p>	

Change History		
Date	Change details, since approval	Approved by

# Paediatric Critical Care

## Drowning

**Drowning:** Respiratory impairment following submersion/immersion in a liquid medium<sup>1</sup>

“Near drowning” and “wet/dry drowning” terms are no longer used.

### PRE-HOSPITAL & CARDIAC ARREST MANAGEMENT

Early bystander CPR dramatically affects chances of survival.<sup>4,5,6</sup>

Emergency medical services (EMS) initiate A, B, C approach

**A:** Immobilise C-Spine. Ensure airway is clear. Administer 100% O<sub>2</sub>.

**B:** Bag mask ventilation/ intubation by skilled personnel if required

**C:** Cardiac compressions 15:2. Obtain IV/IO access if possible.

**If cardiac arrest: use APLS algorithm and adjust if hypothermic<sup>12</sup>:**

- **<30°C:** aggressively rewarm (see right), avoid adrenaline/amiodarone and max 3 defibrillation attempts until >30°C
- **30-35°C:** Defibrillate as usual, double dose interval for resus. drugs

### MANAGEMENT: Discuss with PICU early

#### Protect airway

- Assume cervical injury until proven otherwise
- High flow O<sub>2</sub> (trauma mask)<sup>1,6</sup>
- Intubation with cuffed endotracheal tube.

#### Protective lung strategy

- Target tidal volumes 6-8mls/kg. Limit PIP to 30cmH<sub>2</sub>O
- Permissive hypercapnia. (Caution with head injury see below)
- Permissive hypoxia SaO<sub>2</sub> 90-94%
- Recruitment manoeuvres to establish open lung strategy
- Optimise PEEP to achieve oxygenation, may need 10-15cmH<sub>2</sub>O<sup>10</sup>

#### Circulation<sup>4,6</sup>

- Treat hypothermia to optimise resuscitation process
- (See rewarming methods.)
- Target core temp >35°C
- Central / IO access and arterial access
- Restrict fluids to 50% maintenance (0.9% Sodium Chloride +/- Glucose).
- Age appropriate MAP (inotropes): cerebral perfusion pressure)
- Arrhythmias common and may require treatment. Discuss with STRS

#### Neuroprotection<sup>4,6,8</sup>

- 30 degree head up tilt
- Normoglycaemic
- 2.7% sodium chloride (2 - 5ml/kg over 30mins) target sodium 145- 150mmol/l
- Ventilate to ETCO<sub>2</sub> target 4-5 kPa
- If haemodynamically stable: slow rewarming to 35°C advised<sup>10</sup>
- May need seizure control: load phenytoin

**Full secondary survey**

**Early referral to tertiary centre for definitive care**

### PROGNOSTICATORS OF OUTCOME

#### GOOD

Short submersion time. GCS>5, cardiac output and spontaneous respiration in A&E

#### BAD

Age<3yrs<sup>2,3,5</sup> Submersion> 5 min No CPR>10 mins, asystole at the scene, resuscitation of cardiac arrest > 30mins<sup>9</sup> Multiorgan dysfunction<sup>11</sup>

### REWARMING METHODS

Warm patient with core temp< 28°C<sup>7,8</sup>

If CVS stable aim to rewarm at 0.5°C per hr to 35°C

- Warm IV fluids (38-40°C)
- Heated humidified ventilator gases
- Radiant heaters/ Warming blanket
- Intravascular temperature control devices

**If cardiovascularly unstable/in cardiac arrest:**

Aggressive rewarming

- Bladder irrigation/ pleural lavage warm fluids
- Intravascular temperature control devices
- CVVH
- CPB/ECMO if available.

Prolonged, profound hypothermia may make resuscitation and rewarming impossible.

### INVESTIGATIONS

Blood gas and glucose

FBC, Coagulation

U&E's: Correct any electrolyte abnormalities

CK may be ↑↑ if child has struggled in hypoxic conditions

LFT's & C-reactive protein. Serum osmolality

Consider drug and alcohol screen

Chest x-ray, Trauma panel if indicated

CT Scan/MRI brain if warranted.

### ONGOING MANAGEMENT

- Multi organ dysfunction common in paediatric drowning victims who require PICU (54%).<sup>11</sup>
  - Respiratory>neurologic> cardiovascular
- Maintain normothermia if post cardiac arrest
- Antibiotics not started routinely
  - Send BAL and treat if concerns
- No evidence for the use of corticosteroids<sup>1,2,7,10</sup>
- Minimal data to support the use of surfactant<sup>1</sup>

#### References

1. Robinson, Pediatric Emergency Care 1987 Mar;3(1):44-49.
2. Golden, Br. J. Anaesth 1997;79:214-225.
3. Nichter, Crit Care Med 1989 Oct;17(10):993-995.
4. Burford Pediatric Emergency Care 2005 Sept;21(9):610-616.
5. Quan, J Emerg Nursing 1998 Apr;24(2):1-2.
6. Verive, Medscape Article 2009 Apr;908677:1-3.
7. Lord, J R Army Med Corps 2005;151:250-255.
8. Moon, Emerg Med 2002 Dec;14(4):377-386.
9. Kieboom et al BMJ 2015;350
10. Topijan et al Neurocrit Care 2012; dec 17: 841
11. Mtaweh et al resuscitation 2015 (90): 91
12. APLS 6<sup>th</sup> Edition 2015